

Harvey Mudd College Math Tutorial:

Matrix Algebra

We review here some of the basic definitions and elementary algebraic operations on matrices.

There are many applications as well as much interesting theory revolving around these concepts, which we encourage you to explore after reviewing this tutorial.

A **matrix** is simply a rectangular array of numbers. For example,

$$A = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix}$$

is a $m \times n$ matrix (m rows, n columns), where the entry in the i^{th} row and j^{th} column is a_{ij} . We often write $A = [a_{ij}]$.

Some Terminology

For an $n \times n$ **square** matrix A , the elements $a_{11}, a_{22}, \dots, a_{nn}$ form the **main diagonal** of the matrix. The sum $\sum_{k=1}^n a_{kk}$ of the elements on the main diagonal of A is called the **trace** of A .

The matrix $A^T = [a_{ji}]$ formed by interchanging the rows and columns of A is called the **transpose** of A . If $A^T = A$, the matrix A is **symmetric**.

Example

Let $B = \begin{bmatrix} 6 & 9 \\ -4 & -6 \end{bmatrix}$. The trace of B is $6 + (-6) = 0$.

The transpose of B is $B^T = \begin{bmatrix} 6 & -4 \\ 9 & -6 \end{bmatrix}$.

Addition and Subtraction of Matrices

To **add** or **subtract** two matrices of the same size, simply add or subtract corresponding entries. That is, if $B = [b_{ij}]$ and $C = [c_{ij}]$,

$$B + C = [b_{ij} + c_{ij}] \quad \text{and} \quad B - C = [b_{ij} - c_{ij}].$$

Example

$$\text{For } B = \begin{bmatrix} 6 & 9 \\ -4 & -6 \end{bmatrix} \text{ and } C = \begin{bmatrix} 1 & 2 \\ -1 & 0 \end{bmatrix},$$

$$B + C = \begin{bmatrix} 6+1 & 9+2 \\ -4+(-1) & -6+0 \end{bmatrix} = \begin{bmatrix} 7 & 11 \\ -5 & -6 \end{bmatrix}$$

$$B - C = \begin{bmatrix} 6-1 & 9-2 \\ -4-(-1) & -6-0 \end{bmatrix} = \begin{bmatrix} 5 & 7 \\ -3 & -6 \end{bmatrix}.$$

The $m \times n$ **zero matrix**, $\mathbf{0}$, for which every entry is 0, has the property that for any $m \times n$ matrix A ,

$$A + \mathbf{0} = A.$$

Scalar Multiplication

To multiply a matrix A by a number c (a “scalar”), multiply each entry of A by c . That is,

$$cA = [ca_{ij}].$$

Example

Using the matrix $B = \begin{bmatrix} 6 & 9 \\ -4 & -6 \end{bmatrix}$ from the previous example,

$$3B = 3 \begin{bmatrix} 6 & 9 \\ -4 & -6 \end{bmatrix} = \begin{bmatrix} 18 & 27 \\ -12 & -18 \end{bmatrix}.$$

Matrix Multiplication

Let X be an $m \times n$ matrix and Y be an $n \times p$ matrix. Then the **product** XY is the $m \times p$ matrix whose $(i, j)^{th}$ entry is given by

$$\sum_{k=1}^n x_{ik}y_{kj}.$$

Notes

- The product XY is only defined if the number of columns of X is the same as the number of rows of Y .
- XY and YX may very well not both be defined. If they both do exist, they are not necessarily equal and in fact might not even be of the same size.

Example

For the matrices $B = \begin{bmatrix} 6 & 9 \\ -4 & -6 \end{bmatrix}$ and $C = \begin{bmatrix} 1 & 2 \\ -1 & 0 \end{bmatrix}$,

$$BC = \begin{bmatrix} 6 & 9 \\ -4 & -6 \end{bmatrix} \begin{bmatrix} 1 & 2 \\ -1 & 0 \end{bmatrix} = \begin{bmatrix} (6)(1) + (9)(-1) & (6)(2) + (9)(0) \\ (-4)(1) + (-6)(-1) & (-4)(2) + (-6)(0) \end{bmatrix} = \begin{bmatrix} -3 & 12 \\ 2 & -8 \end{bmatrix}$$

while

$$CB = \begin{bmatrix} 1 & 2 \\ -1 & 0 \end{bmatrix} \begin{bmatrix} 6 & 9 \\ -4 & -6 \end{bmatrix} = \begin{bmatrix} (1)(6) + (2)(-4) & (1)(9) + (2)(-6) \\ (-1)(6) + (0)(-4) & (-1)(9) + (0)(-6) \end{bmatrix} = \begin{bmatrix} -2 & -3 \\ -6 & -9 \end{bmatrix}.$$

The $n \times n$ matrix having all main diagonal entries equal to 1 and all other entries equal to 0 is called the **identity** matrix I . For example, the 3×3 matrix is $\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$. The $n \times n$ identity matrix has the property that if A is any $n \times n$ matrix,

$$AI = IA = A.$$

Inverse of a Matrix

Start with an $n \times n$ matrix X . Suppose the $n \times n$ matrix Y has the property that

$$XY = YX = I.$$

Then Y is called the **inverse** of X and is denoted X^{-1} .

Notes

- Only square matrices X can have inverses. If X is *not* square, then for any Y the product XY will not be the same size matrix as the product YX (if we're lucky enough even to have both products exist!).
- Not every square matrix has an inverse. If an inverse exists, it is unique.
- If a matrix has an inverse, the matrix is said to be **invertible**.

The inverse of a 2×2 matrix is simple to calculate:

$$\text{If } A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}, \text{ then } A^{-1} = \frac{1}{ad - bc} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix}.$$

Example

The inverse of $C = \begin{bmatrix} 1 & 2 \\ -1 & 0 \end{bmatrix}$ is

$$C^{-1} = \frac{1}{(1)(0) - (2)(-1)} \begin{bmatrix} 0 & -2 \\ 1 & 1 \end{bmatrix} = \frac{1}{2} \begin{bmatrix} 0 & -2 \\ 1 & 1 \end{bmatrix} = \begin{bmatrix} 0 & -1 \\ 1/2 & 1/2 \end{bmatrix}.$$

Note that $CC^{-1} = \begin{bmatrix} 1 & 2 \\ -1 & 0 \end{bmatrix} \begin{bmatrix} 0 & -1 \\ 1/2 & 1/2 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$

and $C^{-1}C = \begin{bmatrix} 0 & -1 \\ 1/2 & 1/2 \end{bmatrix} \begin{bmatrix} 1 & 2 \\ -1 & 0 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}.$

Matrix $B = \begin{bmatrix} 6 & 9 \\ -4 & -6 \end{bmatrix}$ does not have an inverse.

Determinant of a Matrix

How did we know that $B = \begin{bmatrix} 6 & 9 \\ -4 & -6 \end{bmatrix}$ does not have an inverse?

The **determinant** of A , $\det A$, is a number with the property that A is invertible if and only if $\det A \neq 0$.

For a 2×2 matrix $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$, $\det A = ad - bc$.

Example

For $B = \begin{bmatrix} 6 & 9 \\ -4 & -6 \end{bmatrix}$, $\det B = (6)(-6) - (9)(-4) = -36 + 36 = 0$, so B is not invertible.

That is, B does not have an inverse.

For a 3×3 (or larger) matrix A , things are a little more complicated:

Denote by $M_{ij}(A)$ the determinant of the matrix formed by deleting row i and column j for A .

Define $c_{ij}(A) = (-1)^{i+j}M_{ij}(A)$ to be the (i, j) **cofactor** of A .

Then we can compute $\det A$ by the **Laplace Expansion** along any row or column of A :

Along row i :

$$\det A = a_{i1}c_{i1}(A) + a_{i2}c_{i2}(A) + \dots + a_{in}c_{in}(A).$$

Along column j :

$$\det A = a_{1j}c_{1j}(A) + a_{2j}c_{2j}(A) + \dots + a_{nj}c_{nj}(A).$$

Example

$$\text{Let } A = \begin{bmatrix} 1 & -1 & 3 \\ 1 & 0 & -1 \\ 2 & 1 & 6 \end{bmatrix}.$$

Along the first row,

$$\begin{aligned} \det A &= (1)[(0)(6) - (-1)(1)] - (-1)[(1)(6) - (-1)(2)] + 3[(1)(1) - (0)(2)] \\ &= (1)(1) + (1)(8) + (3)(1) \\ &= 12. \end{aligned}$$

Computing $\det A$ along the second column instead,

$$\begin{aligned} \det A &= -(-1)[(1)(6) - (-1)(2)] + (0)[(1)(6) - (3)(2)] - 1[(1)(-1) - (3)(1)] \\ &= (1)(8) + (0)(0) - (1)(-4) \\ &= 12 \text{ as expected.} \end{aligned}$$

Key Concepts

Let $A = [a_{ij}]$ and $B = [b_{ij}]$.

- Transpose A^T of A :

$$A^T = [a_{ji}].$$

- Trace of A :

$$\sum_{k=1}^n a_{kk} \text{ (for an } n \times n \text{ matrix } A).$$

- Identity Matrix I :

the $n \times n$ matrix with 1's on the main diagonal and 0's elsewhere.

- $A + B$ and $A - B$:

$$A + B = [a_{ij} + b_{ij}]$$

$$A - B = [a_{ij} - b_{ij}].$$

- Scalar Multiplication:

$$cA = [ca_{ij}].$$

- **Matrix Product AB :**

$(i, j)^{th}$ entry is $\sum_{k=1}^n a_{ik}b_{kj}$

(for an $m \times n$ matrix A and an $n \times p$ matrix B).

- **Inverse A^{-1} of A :**

A^{-1} satisfies $AA^{-1} = A^{-1}A = I$.

If $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$,

then $A^{-1} = \frac{1}{ad-bc} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix}$.

- **Determinant $\det A$:**

If $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$, $\det A = ad - bc$.

In general,

along row i :

$$\det A = a_{i1}c_{i1}(A) + a_{i2}c_{i2}(A) + \dots + a_{in}c_{in}(A).$$

along column j :

$$\det A = a_{1j}c_{1j}(A) + a_{2j}c_{2j}(A) + \dots + a_{nj}c_{nj}(A).$$

[\[I'm ready to take the quiz.\]](#) [\[I need to review more.\]](#)

[\[Take me back to the Tutorial Page\]](#)