

Introduction

A finite continued fraction is an expression of the form

$$a_0 + \frac{b_1}{a_1 + \frac{b_2}{\dots + \frac{b_n}{a_n}}} \quad (1)$$

Although this expression may look messy, there is an elementary, conceptually elegant way to understand continued fractions. In Chapter 3 of their award-winning book [1], Arthur T. Benjamin and Jennifer J. Quinn introduce a clever approach that relies only on counting! This approach works for the well-studied *regular* continued fractions, which have the form

$$a_0 + \frac{1}{\dots + \frac{1}{a_n}} \quad (2)$$

where the a_i are positive integers. The main idea in this chapter is that (2) has a combinatorial counterpart: the set of *square-and-domino tilings* of a length $(n + 1)$ board with height conditions, as shown in Figure 1.




Figure 1: All nine ways to tile the length 3 board that do not exceed the height conditions $a_0 = 1, a_1 = 3, a_2 = 2$.

Research

Our original goal for the project was to relate regular continued fractions to the less-known *negative* continued fractions, which have the form

$$a_0 - \frac{1}{\dots - \frac{1}{a_n}} \quad (3)$$

This meant that we wanted to:

- Prove that the basic properties of regs (for instance, in [2]) hold for negs also
 - Find a combinatorial interpretation for negs
 - Describe how to convert between regs and negs
- Over the course of the year, we accomplished these goals and more.

Results

Weighted Tilings

A continued fraction in general form (1), where a_i and b_i may be any real numbers, requires the concept of a *weighted tiling* to express combinatorially.




Figure 2: All the ways to tile the $|0 : 3|$ board.

We assign a_0, \dots, a_n to the squares and b_1, \dots, b_n to the dominoes, which gives each tile a weight. Then we define $|i : j|$ as the weighted sum of all possible tilings covering the cells from i to j . For instance

$$|0 : 3| = a_0 a_1 a_2 a_3 + a_0 a_1 b_3 + a_0 b_2 a_3 + b_1 a_2 a_3 + b_1 b_3$$

as shown in Figure 2. With this definition, the fundamental connection between continued fractions and combinatorics is given by

$$a_0 + \frac{b_1}{a_1 + \frac{b_2}{\dots + \frac{b_n}{a_n}}} = \frac{|0 : n|}{|1 : n|}$$

Mixed Tilings

For regular continued fractions, the weights aren't needed because having *height conditions* gave us a number of tilings equal the weighted sum $|0 : n|$. This thesis shows how to do the same thing for negs, that is, when $b_i = -1$.




Figure 3: The board representing $1 - \frac{1}{3 - \frac{1}{2}} = \frac{3}{5}$. We use the new objects \rightarrow and \leftarrow instead of dominoes, where $\rightarrow\leftarrow$ cannot appear consecutively.

Converting between Regs and Negs

Given a sequence a_0, a_1, \dots, a_n for a regular continued fraction,

- Increase the even-indexed numbers a_{2k} by 2 (except the one(s) on the ends, which are increased by 1), and
- Replace the odd-indexed numbers a_{2k+1} by a sequence of $(a_{2k+1} - 1) 2s$

in order to obtain the negative continued fraction for the same number. For instance,

$$6 + \frac{1}{3 + \frac{1}{3}} = 7 - \frac{1}{2 - \frac{1}{2 - \frac{1}{4}}}$$

Although this pattern is hard to guess, it's not hard to see combinatorially!

An Uncounted Identity

Another application of this framework is a new proof of the following identity:

Identity 1 Let F_n denote the n th Fibonacci number and L_n denote the n th Lucas number. Then, for all $m, n \in \mathbb{Z}^+$,

$$\frac{F_{(n+1)m}}{F_{nm}} = L_m - \frac{(-1)^m}{L_m - \frac{(-1)^m}{\dots - \frac{(-1)^m}{L_m}}}$$

where the term L_m appears n times in the continued fraction.

In fact, by generalizing the Fibonacci and Lucas numbers, we obtain an interesting theorem about periodic continued fractions!

Conclusions

We focused on results and identities involving negative continued fractions, but a number of other topics related to continued fractions could surely be addressed using similar methods, such as,

- Generalizations of other Fibonacci identities
- Periodic continued fractions
- Two-squares theorems (from number theory)
- Class numbers of quadratic number fields.

It's always interesting to see the complex results that "fall out" as simple consequences of the combinatorial approach. Prof. Benjamin and I will be excited to see more such results from future thesis students! Moreover, only high school mathematics is really required to really dig into these subjects, making them ideal for educational purposes.

References

1. Arthur T. Benjamin and Jennifer J. Quinn. *Proofs That Really Count: The Art of Combinatorial Proof*, Mathematical Association of America, Washington DC, 2003.
2. Andrew M. Rockett and Peter Szűsz. *Continued fractions*. World Scientific Publishing Co., New York, 1964.

Acknowledgments

My advisor, Professor Benjamin, made this topic possible in the first place and provided excellent suggestions and guidance. I'd also like to thank Sanjai Gupta, my second reader; Professor Darryl Yong for helpful \LaTeX tips; and Professor Richard Guy for suggesting some research ideas.

For Further Information

To download the thesis or this poster, please visit <http://www.math.hmc.edu/~aeustis/thesis/>. To contact us via email:

- Alex Eustis: akeustis@gmail.com
- Arthur T. Benjamin: benjamin@math.hmc.edu