

# A COMBINATORIAL APPROACH TO FIBONOMIAL COEFFICIENTS

**Arthur T. Benjamin and Sean S. Plott**

Dept. of Mathematics, Harvey Mudd College, Claremont, CA 91711  
benjamin@hmc.edu

ABSTRACT. A combinatorial argument is used to explain the integrality of Fibonomial coefficients and their generalizations. The numerator of the Fibonomial coefficient counts tilings of staggered lengths, which can be decomposed into a sum of integers, such that each integer is a multiple of the denominator of the Fibonomial coefficient. By colorizing this argument, we can extend this result from Fibonacci numbers to arbitrary Lucas sequences.

## 1. INTRODUCTION

The Fibonomial Coefficient  $\binom{n}{k}_F$  is defined, for  $0 < k \leq n$ , by replacing each integer appearing in the numerator and denominator of  $\binom{n}{k} = \frac{n(n-1)\cdots(n-k+1)}{k(k-1)\cdots 1}$  with its respective Fibonacci number. That is,

$$\binom{n}{k}_F = \frac{F_n F_{n-1} \cdots F_{n-k+1}}{F_k F_{k-1} \cdots F_1}.$$

For example,  $\binom{7}{3}_F = \frac{F_7 F_6 F_5}{F_3 F_2 F_1} = \frac{13 \cdot 8 \cdot 5}{2 \cdot 1 \cdot 1} = 260$ .

It is, at first, surprising that this quantity will always take on integer values. This can be shown by an induction argument by replacing  $F_n$  in the numerator with  $F_k F_{n-k+1} + F_{k-1} F_{n-k}$ , resulting in

$$\binom{n}{k}_F = F_{n-k+1} \binom{n-1}{k-1}_F + F_{k-1} \binom{n-1}{k}_F.$$

By similar reasoning, this integrality property holds for any Lucas sequence defined by  $U_0 = 0$ ,  $U_1 = a$  and for  $n \geq 2$ ,  $U_n = aU_{n-1} + bU_{n-2}$ , and we define

$$\binom{n}{k}_U = \frac{U_n U_{n-1} \cdots U_{n-k+1}}{U_k U_{k-1} \cdots U_1}.$$

In this note, we combinatorially explain the integrality of  $\binom{n}{k}_F$  and  $\binom{n}{k}_U$  by a tiling interpretation, answering a question proposed in Benjamin and Quinn's book, *Proofs That Really Count* [1].

## 2. STAGGERED TILINGS

It is well known that for  $n \geq 0$ ,  $f_n = F_{n+1}$  counts tilings of a  $1 \times n$  board with squares and dominoes [1]. For example,  $f_4 = 5$  counts the five tilings of length four, where  $s$  denotes a square tile and  $d$  denotes a domino tile:  $ssss, ssd, sds, dss, dd$ . Hence, for  $\binom{n}{k}_F = \frac{f_{n-1}f_{n-2}\cdots f_{n-k}}{f_{k-1}f_{k-2}\cdots f_0}$ , the numerator counts the ways to simultaneously tile boards of length  $n-1, n-2, \dots, n-k$ . The challenge is to find disjoint “subtilings” of lengths  $k-1, k-2, \dots, 0$  that can be described in a precise way. Suppose  $T_1, T_2, \dots, T_k$  are tilings with respective lengths  $n-1, n-2, \dots, n-k$ . We begin by looking for a tiling of length  $k-1$ .

If  $T_1$  is “breakable” at cell  $k-1$ , which can happen  $f_{k-1}f_{n-k}$  ways, then we have found a tiling of length  $k-1$ . We would then look for a tiling of length  $k-2$ , starting with tiling  $T_2$ .

Otherwise,  $T_1$  is breakable at cell  $k-2$ , followed by a domino (which happens  $f_{k-2}f_{n-k-1}$  ways. Here, we “throw away” cells 1 through  $k$ , and consider the remaining cells to be a new tiling, which we call  $T_{k+1}$ . (Note that  $T_{k+1}$  has length  $n-k-1$ , which is one less than the length of  $T_k$ .) We would then continue our search for a tiling of length  $k-1$  in  $T_2$ , then  $T_3$ , and so on, creating  $T_{k+2}, T_{k+3}$ , and so on as we go, until we eventually find a tiling  $T_{x_1}$  that is breakable at cell  $k-1$ . (We are guaranteed that  $x_1 \leq n-k+1$  since  $T_{n-k+1}$  has length  $k-1$ .) At this point, we disregard everything in  $T_{x_1}$  and look for a tiling of length  $k-2$ , beginning with tiling  $T_{x_1+1}$ .

Following this procedure, we have, for  $1 \leq x_1 < x_2 < \cdots < x_{k-1} \leq n$ , the number of tilings  $T_1, T_2, \dots, T_k$  that lead to finding a tiling of length  $k-i$  at the beginning of tiling  $T_{x_i}$  is

$$f_{k-2}^{x_1-1} f_{k-1} f_{n-x_1-(k-1)} f_{k-3}^{x_2-x_1-1} f_{k-2} f_{n-x_2-(k-2)} \cdots f_0^{x_{k-1}-x_{k-2}-1} f_1 f_{n-x_{k-1}-1}.$$

Consequently, if we define  $x_0 = 0$ , then  $F_n F_{n-1} \cdots F_{n-k+1}$

$$\begin{aligned} &= f_{n-1} f_{n-2} \cdots f_{n-k} \\ &= f_{k-1} f_{k-2} f_{k-3} \cdots f_1 \sum_{1 \leq x_1 < x_2 < \cdots < x_{k-1} \leq n-1} \cdots \sum_{i=1}^{k-1} \prod_{i=1}^{k-1} (f_{k-1-i})^{x_i-x_{i-1}-1} f_{n-x_i-(k-i)} \\ &= F_k F_{k-1} F_{k-2} \cdots F_2 F_1 \sum_{1 \leq x_1 < x_2 < \cdots < x_{k-1} \leq n-1} \cdots \sum_{i=1}^{k-1} \prod_{i=1}^{k-1} (F_{k-i})^{x_i-x_{i-1}-1} F_{n-x_i-(k-i)+1}. \end{aligned}$$

That is,

$$\binom{n}{k}_F = \sum_{1 \leq x_1 < x_2 < \cdots < x_{k-1} \leq n-1} \cdots \sum_{i=1}^{k-1} \prod_{i=1}^{k-1} F_{k-i}^{x_i-x_{i-1}-1} F_{n-x_i-(k-i)+1}.$$

This theorem has a natural Lucas sequence generalization. For positive integers  $a, b$ , it is shown in [1] that  $u_n = U_{n+1}$  counts colored tilings of length  $n$ , where

there are  $a$  colors of squares and  $b$  colors of dominoes. (More generally, if  $a$  and  $b$  are any complex numbers,  $u_n$  counts the total weight of length  $n$  tilings, where squares and dominoes have respective weights  $a$  and  $b$ , and the weight of a tiling is the product of the weights of its tiles.) By virtually the same argument as before, we have

$$\binom{n}{k}_U = \sum_{1 \leq x_1 < x_2 < \dots < x_{k-1} \leq n-1} \prod_{i=1}^{k-1} b^{x_{k-1} - (k-1)} U_{k-i}^{x_i - x_{i-1} - 1} U_{n-x_i - (k-i) + 1}.$$

The presence of the  $b^{x_{k-1} - (k-1)}$  term accounts for the  $x_{k-1} - (k-1)$  dominoes that caused  $x_{k-1} - (k-1)$  tilings to be unbreakable at their desired spot.

As an immediate corollary, we note that the right hand side of this identity is a multiple of  $b$ , unless  $x_i = i$  for  $i = 1, 2, \dots, k-1$ . It follows that

$$\binom{n}{k}_U \equiv U_{n-k+1}^{k-1} \pmod{b}.$$

#### REFERENCES

- [1] A. T. Benjamin and J. J. Quinn, *Proofs That Really Count: The Art of Combinatorial Proof*, Washington DC: Mathematical Association of America, 2003.

AMS Subject Classification Numbers: 05A19, 11B39.